


12.4 Applications: Students should review 7.5

Center of Mass




balances $\Leftrightarrow m_1 d_1 = m_2 d_2$

Law of the Lever

Archimedes ~ 200 B.C.

Suppose we have



How do we find the balance pt \bar{x} ?

Let $d_1 = \bar{x} - x_1$, $d_2 = x_2 - \bar{x}$.


Then $m_1(\bar{x} - x_1) = m_2(x_2 - \bar{x}) \Rightarrow \bar{x} = \frac{m_1 x_1 + m_2 x_2}{m_1 + m_2}$.

If there are several weights

$$\bar{x} = \frac{m_1 x_1 + m_2 x_2 + \dots + m_n x_n}{m_1 + \dots + m_n} = \frac{\sum_{i=1}^n m_i x_i}{\sum_{i=1}^n m_i}$$

weighted average
& total mass

Now let $p(x)$ be a linear density function, $p(x) = \frac{\text{mass}}{\text{length}}$.



Partition $[a, b]$ by $a = x_0 < x_1 < \dots < x_n = b$.

Let $m_i = p(x_i) \Delta x$.


Then
$$\bar{x} \approx \frac{\sum m_i x_i}{\sum m_i} = \frac{\sum p(x_i) x_i \Delta x}{\sum p(x_i) \Delta x}$$

Take the limits as $n \rightarrow \infty, \Delta x \rightarrow 0$ to get

$$\bar{x} = \frac{\int_a^b p(x) x dx}{\int_a^b p(x) dx}$$



Ex Let $p(x) = x^2$ over $[0, 1]$. Find balance pt, \bar{x} .

$$\bar{x} = \frac{\int_0^1 x^3 dx}{\int_0^1 x^2 dx} = \frac{1/4}{1/3} = \frac{3}{4}$$



Now we want to generalize this to 2-dimensions

Let $\rho(x,y)$ be a density over a rectangle R ; $\rho(x,y) = \frac{\text{mass}}{\text{area}}$
 Let R be $[a, b] \times [c, d]$. Partition as below



Let $m_{ij} = \rho(x_i, y_j) \Delta x \Delta y \approx$ mass in cell ij .

Let's find \bar{x} . Let $M_i = \sum_{j=1}^m m_{ij} \approx$ mass of column i .
 Then

$$\bar{x} \approx \frac{\sum M_i x_i}{\sum M_i} = \frac{\sum \sum x_i \rho(x_i, y_j) \Delta x \Delta y}{\sum \sum \rho(x_i, y_j) \Delta x \Delta y}$$


Take limits to get

$$\bar{x} = \frac{\int_a^b \int_c^d x \rho(x,y) dx dy}{\iint \rho(x,y) dx dy} \leftarrow \text{mass } M$$

Like wise

$$\bar{y} = \frac{M_x}{M} = \frac{\int \int y \rho(x,y) dx dy}{\iint \rho(x,y) dx dy}$$

Ex Find the mass and center of mass of the square $[0,1] \times [0,1]$, with density $\rho(x,y) = x^2 + 2y + 1$



$$M = \int_0^1 \int_0^1 (x^2 + 2y + 1) dx dy = \int_0^1 \left(\frac{1}{3} + 2y + 1 \right) dy = \frac{4}{3} + 1 = \frac{7}{3}$$

$$M_y = \int_0^1 \int_0^1 x(x^2 + 2y + 1) dx dy = \frac{5}{4}$$


$$M_x = \int_0^1 \int_0^1 y(x^2 + 2y + 1) dx dy = \frac{4}{3}$$

$$\bar{x} = \frac{M_y}{M} = \frac{5/4}{7/3} = \frac{15}{28} \approx 0.5357142$$

$$\bar{y} = \frac{M_x}{M} = \frac{4/3}{7/3} = \frac{4}{7} = \frac{16}{28} \approx 0.571428571$$

Ex

Let R be the triangular region below. Let $\rho(x,y) = xy$ be a density function. Find the mass and center of mass.



$$\text{Mass} = \int_0^1 \int_y^{2-y} xy \, dx \, dy = \int_0^1 \frac{1}{2} x^2 y \Big|_y^{2-y} dy$$

$$\frac{1}{2} \int_0^1 (2-y)^2 y - y^3 \, dy = \frac{1}{2} \int_0^1 4y - 4y^2 + y^3 - y^3 \, dy$$

$$= 2 \int_0^1 y - y^2 \, dy = 2 \left(\frac{1}{2} - \frac{1}{3} \right) = \frac{2}{6} = \boxed{\frac{1}{3}}$$

$$M_x = \int_0^1 \int_y^{2-y} xy^2 \, dx \, dy = \dots = 2 \int_0^1 y^2 - y^3 \, dy = 2 \left(\frac{1}{3} - \frac{1}{4} \right) = \frac{2}{12} = \boxed{\frac{1}{6}}$$

$$M_y = \int_0^1 \int_y^{2-y} x^2 y \, dx \, dy = \int_0^1 \frac{1}{3} x^3 y \Big|_y^{2-y} dy =$$


$$\frac{1}{3} \int_0^1 (2-y)^3 y - y^4 \, dy = \frac{1}{3} \int_0^1 [8 - 12y + 6y^2 - y^3] y - y^4 \, dy$$

$$= \frac{1}{3} \int_0^1 8y - 12y^2 + 6y^3 - 2y^4 \, dy = \frac{1}{3} \left(\frac{8}{2} - \frac{12}{3} + \frac{6}{4} - \frac{2}{5} \right)$$

$$= \frac{1}{3} \left(\frac{7}{2} - \frac{2}{5} \right) = \frac{1}{3} \left(\frac{15-4}{10} \right) = \boxed{\frac{11}{30}}$$


$$\bar{x} = \frac{M_y}{M} = \frac{\frac{11}{30}}{\frac{1}{3}} = \frac{11}{10} = 1.1$$

$$\bar{y} = \frac{M_x}{M} = \frac{\frac{1}{6}}{\frac{1}{3}} = \frac{1}{2} = 0.5$$



Rotational Inertia

Idea:




$$\omega = \frac{d\theta}{dt}$$

arc length is \$r\theta\$.

$$v = r\omega$$


$$E = \frac{1}{2} m v^2 = \frac{1}{2} m r^2 \omega^2$$

↓
\$I_0\$ or \$I_z\$.



(Same \$\omega\$).

$$\begin{aligned} E &= \sum \frac{1}{2} m_i v_i^2 \\ &= \sum \frac{1}{2} m_i r_i^2 \omega^2 = \\ &= \frac{1}{2} (\sum m_i r_i^2) \omega^2 \\ &= \frac{1}{2} I_0 \omega^2 \end{aligned}$$



$$E \approx \frac{1}{2} \left(\sum \Delta m_i r_i^2 \right) \omega^2 \Rightarrow \frac{1}{2} \left(\sum r_i^2 \rho(x_i, y_i) \Delta x \Delta y \right) \omega^2$$

$$I_0 = \iint_R r^2 \rho \, dA$$

$$E = \text{limit} = \frac{1}{2} \left(\iint_R r^2 \rho(x, y) \, dx \, dy \right) \omega^2$$

$$\text{or } = \frac{1}{2} \left(\iint \rho(r, \theta) r^3 \, dr \, d\theta \right) \omega^2$$

Notice, $I_0 = \iint (x^2 + y^2) \rho \, dx \, dy =$

$$\iint x^2 \rho \, dx \, dy + \iint y^2 \rho \, dx \, dy -$$

$$\parallel$$

$$I_y$$


$$\parallel$$

$$I_x$$

$$I_z = I_0 = I_x + I_y,$$

Ex Find I_0 for the region semi-circle with radius R and density proportional to the distance from the diameter, where I_0 means w.r.t the circle's center.


Sol.



$$\rho(x, y) = ky, \text{ for some } k > 0.$$

$$I = \int_0^\pi \int_0^R k r \sin \theta \, r^3 \, dr \, d\theta = k \int_0^\pi \frac{R^5}{5} \sin \theta \, d\theta =$$

$$\frac{2}{5} k R^5.$$



Ex Find I_0 for the rectangle $[0, 2] \times [0, 1]$ with $p(x, y) = 3x + 2y + 1$.

Sol: $I_x = \int_0^1 \int_0^2 y^2 (3x + 2y + 1) dx dy$

$$= \int_0^1 \int_0^2 3xy^2 + 2y^3 + y^2 dx dy$$

$$\int_0^1 \left. \frac{3x^2 y^2}{2} + (2y^3 + y^2)x \right|_0^2 dy$$

$$= \int_0^1 6y^2 + 4y^3 + 2y^2 dy$$

$$= \left. \frac{8y^3}{3} + y^4 \right|_0^1 = \frac{8}{3} + 1 = \frac{11}{3} = 3\frac{2}{3}$$

$$I_y = \int_0^1 \int_0^2 x^2 (3x + 2y + 1) dx dy$$

$$= \int_0^1 \int_0^2 3x^3 + 2x^2 y + x^2 dx dy =$$

$$= \int_0^1 \left(\frac{3 \cdot 16}{4} + \frac{2 \cdot 8 y}{3} + \frac{8}{3} \right) dy = \int_0^1 \frac{16}{3} y + \frac{4y}{3} dy$$


$$= \frac{8}{3} + \frac{44}{3} = \frac{52}{3} = 17\frac{1}{3}$$

$$I_0 = 3\frac{2}{3} + 17\frac{1}{3} = 21.$$

$$E = \frac{1}{2} I_0 w^2$$

Radius of Gyration.

$$\text{Let } R_G = \sqrt{\frac{I_0}{M}}$$



$$I = MR_G^2 = I_0$$


Ex The radius of Gy of the last example is,

$$R_G = \sqrt{\frac{I_0}{M}}$$

$$M = \int_0^1 \int_0^2 (3x + 2y + 1) dx dy =$$

$$= \int_0^1 (6 + 4y + 2) dy = 8 + 2 = 10$$

$$R_G = \sqrt{\frac{21}{10}} \approx 1.449137675\dots$$



~~$$I = \frac{1}{2} \cdot (0.72)^2 \cdot 10 = 10.0$$~~

Ex

A disk with radius 1 meter and mass 3 kg is moving 10 m/s in a straight line and spinning 5 revolutions per second. It has uniform density. Find the total kinetic energy.

Sol

$E = \frac{1}{2}mv^2 + \frac{1}{2}I_0\omega^2$. We know m and v , and $\omega = 5 \frac{\text{rev}}{\text{sec}} = 5 \frac{\text{rev}}{\text{sec}} \cdot 2\pi \frac{\text{radians}}{\text{rev}} = 10\pi \frac{\text{radians}}{\text{sec}}$. We just need to find I_0 .

$$I_0 = \int_0^{2\pi} \int_0^1 r^2 \rho r dr d\theta. \quad \rho = \frac{\text{mass}}{\text{area}} = \frac{3 \text{ kg}}{\pi \text{ m}^2}$$

$$I_0 = \frac{3}{\pi} \int_0^{2\pi} \int_0^1 r^3 dr d\theta = \frac{3}{\pi} \frac{1}{4} \cdot 2\pi = \frac{3}{2}.$$

$$\text{Thus, } E = \frac{1}{2} \cdot 3(10)^2 + \frac{1}{2} \left(\frac{3}{2}\right) (10\pi)^2 = 150 + 75\pi^2$$


$$\approx 896.22 \text{ J.}$$

↑
Joules

$$= \frac{\text{kg m}^2}{\text{s}^2}$$

Additional Examples if Time Permits

Ex Let R be the square $[0, 1] \times [0, 1]$. Let $\rho(x, y) = xy + 1$. Find the mass, the center of mass, the rotational inertia wrt to $(0, 0)$ and the radius of gyration.



Sol $M = \int_0^1 \int_0^1 xy + 1 \, dx \, dy = \int_0^1 \frac{1}{2}y + 1 \, dy = \frac{1}{2} + 1 = \frac{3}{2}$.

$$M_x = \int_0^1 \int_0^1 y(xy + 1) \, dx \, dy = \int_0^1 \int_0^1 xy^2 + y \, dx \, dy = \int_0^1 \frac{1}{2}y^2 + y \, dy$$
$$= \frac{1}{6} + \frac{1}{2} = \frac{2}{3}$$

$M_y = M_x$ by symmetry. Look for short cuts like this!

Thus $\bar{x} = \bar{y} = \frac{2/3}{3/2} = \frac{2}{3} \cdot \frac{2}{3} = \frac{4}{9}$. The C.M. = $(\frac{4}{9}, \frac{4}{9})$.
↑ center of mass.

$$I_x = \int_0^1 \int_0^1 y^2(xy + 1) \, dx \, dy = \int_0^1 \int_0^1 xy^3 + y^2 \, dx \, dy$$
$$= \int_0^1 \frac{1}{2}y^3 + y^2 \, dy = \frac{1}{2} \cdot \frac{1}{4} + \frac{1}{3} = \frac{11}{24}$$


$I_y = I_x$ by symmetry. Thus, $I_0 = \frac{11}{24} + \frac{11}{24} = \frac{22}{24} = \frac{11}{12}$.

$$R_G = \sqrt{\frac{I_0}{M}} = \sqrt{\frac{11/12}{3/2}} = \sqrt{\frac{11}{18}} \approx 0.8563$$

Done!

Ex Let R be the region determined by $0 \leq r \leq 1 + \cos \theta$. The density is proportional to the distance from the x -axis. Find M , \bar{x} , \bar{y} , I_0 , and R_0 .

Sol Graph the region.



It is a cardioid - a heart shape.

Find the density function. $\rho(x, y) = k|y| = kr|\sin \theta|$.
↑ for some k

Notice both the region and the density are symmetric across the x -axis.

$$M = \int_0^{2\pi} \int_0^{1+\cos\theta} kr|\sin\theta| r dr d\theta = k \int_0^{2\pi} \int_0^{1+\cos\theta} r^2 |\sin\theta| dr d\theta$$

$$= 2k \int_0^{\pi} \int_0^{1+\cos(\theta)} r^2 \sin\theta dr d\theta = 2k \int_0^{\pi} \frac{r^3}{3} \sin\theta \Big|_0^{1+\cos\theta} d\theta$$


$$= \frac{2k}{3} \int_0^{\pi} (1+\cos\theta)^3 \sin\theta d\theta.$$

Let $u = 1 + \cos \theta$. Then $du = -\sin \theta d\theta$ and u goes from 2 to 0. Thus,

$$M = \frac{-2k}{3} \int_2^0 u^3 du = \frac{2k}{3} \int_0^2 u^3 du = \frac{2k}{3} \frac{16}{4} = \frac{8}{3} k.$$

$$M_x = \int_0^{2\pi} \int_0^{1+\cos\theta} \underbrace{(r \sin\theta)}_y k r |\sin\theta| r dr d\theta = \boxed{0}$$

Why? This integrand is odd w.r.t. y .



$$M_y = \int_0^{2\pi} \int_0^{1+\cos\theta} \underbrace{(r \cos\theta)}_x k r |\sin\theta| r dr d\theta$$

$$= 2k \int_0^{\pi} \int_0^{1+\cos\theta} r^3 \cos\theta \sin\theta dr d\theta$$

$$= \frac{2k}{4} \int_0^{\pi} (1+\cos\theta)^4 \cos\theta \sin\theta d\theta$$

Let $u = 1 + \cos\theta$. Then $du = -\sin\theta d\theta$ and u goes from 2 to 0. Also notice $\cos\theta = u - 1$. Thus

$$M_y = -\frac{k}{2} \int_2^0 u^4 (u-1) du = \frac{k}{2} \int_0^2 u^5 - u du$$

$$= \frac{k}{2} \left(\frac{2^6}{6} - \frac{2^2}{2} \right) = k \left(\frac{2^4}{3} - 1 \right) = k \left(5\frac{1}{3} - 1 \right)$$

$$= \boxed{\frac{13}{3} k}$$

~~C.M. is (0, 0)~~

$$\bar{x} = \frac{0}{M} = 0. \quad \bar{y} = \frac{M_x}{M} = \frac{\frac{13}{3}k}{\frac{8}{3}k} = \frac{13}{8}$$

$$\text{C.M.} = \left(0, \frac{13}{8}\right).$$

↳ We know by symmetry.

$$I_0 = k \int_0^{2\pi} \int_0^{1+\cos\theta} r^2 r \sin\theta \, r \, dr \, d\theta =$$

$$2k \int_0^{\pi} \int_0^{1+\cos\theta} r^4 \sin\theta \, dr \, d\theta =$$

$$\frac{2k}{5} \int_0^{\pi} (1+\cos\theta)^5 \sin\theta \, d\theta.$$

Let $u = 1 + \cos\theta$. Then $du = -\sin\theta \, d\theta$, $u = 2$ to 0 .

$$I_0 = -\frac{2k}{5} \int_2^0 u^5 \, du = \frac{2}{5}k \int_0^2 u^5 \, du = \frac{2}{5}k \left. \frac{u^6}{6} \right|_0^2 = \boxed{\frac{64}{15}k}.$$

$$R_G = \sqrt{\frac{I_0}{M}} = \sqrt{\frac{\frac{64}{15}k}{\frac{8}{3}k}} = \sqrt{\frac{16}{5}} = \frac{4}{\sqrt{5}} \approx \boxed{1.78885}$$

Done!