



## Section 8 Cones

Def

Let  $K$  be an  $n$ -dim. complex in  $\mathbb{R}^m$ ,  $m > n$ .  
Let  $w \in \mathbb{R}^m$  be s.t. every ray from  $w$  meets  $|K|$   
in at most one point. Let

$$w * K = \{ [w, a_0, \dots, a_p] \mid [a_0, \dots, a_p] \in K \} \cup \{K\} \cup \{w\}.$$

Then  $w * K$  is an  $n+1$ -dim. complex called the  
cone with base  $K$  and vertex  $w$ ,



Notation

For  $\sigma = [a_0, \dots, a_p]$ , let  $[w, \sigma] = [w, a_0, \dots, a_p]$ .

If  $c = \sum n_i \sigma_i$  is a  $p$ -chain, let  $[w, c] = \sum n_i [w, \sigma_i]$ .

This bracket operation,

$$[w, -] : C_p(K) \rightarrow C_{p+1}(w * K)$$

is a group homomorphism.

Lemma

$$(1) \quad \partial [w, \sigma] = \begin{cases} \sigma - w & \text{if } \dim \sigma = 0 \\ \sigma - [w, \partial \sigma] & \text{if } \dim \sigma > 0. \end{cases}$$

(2) Let  $c_0 \in C_0(K)$  and  $c_p \in C_p(K)$ . Then

$$\partial [w, c_0] = c_0 - \varepsilon(c_0)w \quad (\varepsilon(\sum n_i v_i) = \sum n_i)$$

$$\partial [w, c_p] = c_p - [w, \partial c_p], \quad p > 0.$$

Pf

If  $\dim \sigma = 0$ , then  $\sigma$  is a vertex, so  $[\omega, \sigma]$  is an edge. Hence  $\partial[\omega, \sigma] = \sigma - \omega$ .

$$\text{Let } c_0 = \sum n_i v_i. \text{ Then } \partial[\omega, c_0] = \partial \sum n_i [\omega, v_i]$$

$$= \sum n_i \partial[\omega, v_i] = \sum n_i (v_i - \omega) = \sum n_i v_i - \sum n_i \omega$$

$$= c_0 - \varepsilon(c_0) \omega.$$

$$\text{Let } \sigma = [a_0, \dots, a_p], \quad p > 0. \text{ Then } \partial[\omega, \sigma] = \partial[\omega, a_0, \dots, a_p]$$

$$= [a_0, \dots, a_p] + \sum_{i=0}^p (-1)^{i+1} [\omega, a_0, \dots, \hat{a}_i, \dots, a_p]$$

$$= \sigma - [\omega, \sum_{i=0}^p (-1)^i [a_0, \dots, \hat{a}_i, \dots, a_p]] = \sigma - [\omega, \partial \sigma].$$

$$\text{Let } c_p = \sum n_i \sigma_i \in C_p(k), \quad p > 0. \text{ Then}$$

$$\partial[\omega, c_p] = \partial \left( \sum n_i [\omega, \sigma_i] \right) = \sum n_i \partial[\omega, \sigma_i]$$



$$= \sum n_i (\sigma_i - [\omega, \partial \sigma_i]) = \sum n_i \sigma_i - \sum n_i [\omega, \partial \sigma_i]$$

$$= c_p - [\omega, \sum n_i \partial \sigma_i] = c_p - [\omega, \partial \sum n_i \sigma_i]$$

$$= c_p - [\omega, \partial c_p].$$



Ex




$$\text{Let } c_1 = e_1 + e_2.$$

$$\partial c_1 = v_2 - v_0.$$

$$\partial [W, c_1] = c_1 - [W, \partial c_1] = c_1 - [W, v_2] + [W, v_0]$$

$$= e_1 + e_2 + [v_2, w] + [w, v_0]$$



Thm 8.2  $\tilde{H}_p(W * K) = 0, \forall p.$  [We say  $W * K$  is acyclic when this happens.]

Pf The proof is just a calculation using the lemma. It is not deep, but is good practice.

$\tilde{H}_0(W * K) = 0$  since  $|W * K|$  is connected even if  $|K|$  is not.

Let  $p > 0$ . We claim  $H_p(W * K) = 0$ . Let  $z_p \in Z_p(W * K)$ . We will find a  $p+1$ -chain in  $W * K$  with boundary  $z_p$ .

If  $z_p$  is carried by  $K$  then  $\partial [W, z_p] = z_p - [W, \partial z_p] = z_p - [W, 0] = z_p$  and we are done. If not, let  $c_p$  be the part of  $z_p$  carried by  $K$ . We claim,  $z_p = \partial [W, c_p]$ . Before proving this observe that in the example above

$$e_1 + e_2 + [v_2, w] + [w, v_0] = \partial [W, e_1 + e_2].$$

For each simplex  $\sigma$  of  $Z_p - C_p$ ,  $\exists$  a  $p-1$ -simplex of  $K$ ,  $\sigma'$ , with  $\sigma = [w, \sigma']$  by definition. Thus,

$$\exists d_{p-1} \in C_{p-1}(K) \text{ s.t. } Z_p - C_p = [w, d_{p-1}].$$

$$\begin{aligned} \text{Now } Z_p - \partial[w, c_p] &= C_p + [w, d_{p-1}] - (C_p - [w, \partial c_p]) \\ &= [w, d_{p-1}] + [w, \partial c_p] = [w, d_{p-1} + \partial c_p]. \end{aligned}$$

$$\text{Let } e_{p-1} = d_{p-1} + \partial c_p. \text{ Thus } Z_p - \partial[w, c_p] = [w, e_{p-1}]. \quad (\#)$$

$$\text{Next } \partial[w, e_{p-1}] = \partial Z_p - \partial \partial[w, c_p] = 0 - 0 = 0.$$

By the lemma,

$$\left. \begin{array}{l} \text{if } p=1, \quad e_{p-1} - \varepsilon(e_{p-1})w = 0, \text{ and} \\ \text{if } p>1, \quad e_{p-1} - [w, \partial e_{p-1}] = 0. \end{array} \right\} (\#)$$

But  $e_{p-1}$  is carried by  $K$  and  $w$  and no part of  $[w, \partial e_{p-1}]$  is carried by  $K$ . Thus, the terms in  $(\#)$  are independent. Thus,  $e_{p-1} = 0$ .

$$\text{But } (\#) \Rightarrow Z_p - \partial[w, c_p] = [w, 0] = 0 \Rightarrow Z_p = \partial[w, c_p].$$

Thus,  $Z_p \in B_p$  and so  $H_p = Z_p / B_p = 0, p > 1$ .



Thm 8.3 Let  $\sigma$  be an  $n$ -simplex.

(a) Let  $K_\sigma$  be the complex ~~consisting~~ consisting of  $\sigma$  and all faces of  $\sigma$ . Then  $K_\sigma$  is acyclic.

(b) Assume  $n \geq 1$ . Let  $\Sigma_\sigma$  be the complex of all faces of  $\sigma$ . Then

$$\tilde{H}_p(\Sigma_\sigma) \cong \begin{cases} \mathbb{Z} & p = n-1 \\ 0 & \text{otherwise} \end{cases}$$

In the  $p = n-1$  case,  $\tilde{H}_p(\Sigma_\sigma)$  is generated by  $\partial\sigma$ .  
(Assume  $\sigma$  has been given an orientation.)

pf (a) If  $n=0$ ,  $|K_\sigma|$  is a point and so  $\tilde{H}_0(K_\sigma) = 0$ .  
For  $n \geq 1$ ,  $K_\sigma$  is a cone using any vertex for  $w$  and the opposite face for the base. We know cones are acyclic.

(b) If  $n=1$ , then  $K_\sigma$  is an edge with its end points. Then  $|\Sigma_\sigma|$  is two points so  $\tilde{H}_0(\Sigma_\sigma) \cong \mathbb{Z}$ .

Suppose  $n > 1$ . We study the chain groups for  $K_\sigma$  and  $\Sigma_\sigma$ . They are the same except for the top dimension,  $n$ .

$$\begin{array}{ccccccc}
 \langle \sigma \rangle & & & & & & \\
 \parallel & & & & & & \\
 C_n(K_\sigma) & \xrightarrow{d_n} & C_{n-1}(K_\sigma) & \xrightarrow{d_{n-1}} & C_{n-2}(K_\sigma) & \xrightarrow{d_{n-2}} & \dots \xrightarrow{d_1} C_0(K_\sigma) \xrightarrow{\varepsilon} \mathbb{Z} \\
 & & \parallel & & \parallel & & \parallel \\
 C_n(\Sigma_\sigma) & \xrightarrow{d'_n} & C_{n-1}(\Sigma_\sigma) & \xrightarrow{d'_{n-1}} & C_{n-2}(\Sigma_\sigma) & \rightarrow \dots \rightarrow & C_0(\Sigma_\sigma) \xrightarrow{\varepsilon'} \mathbb{Z} \\
 \parallel & & & & & & \parallel \\
 0 & & & & & & 0
 \end{array}$$

The images and kernels of  $d_p$  and  $d'_p$  are the same for  $p \leq n-2$ . Hence  $\tilde{H}_p(\Sigma_\sigma) = \tilde{H}_p(K_\sigma) = 0$  for  $p \leq n-2$ .

Consider  $\tilde{H}_{n-1}(\Sigma_\sigma)$ . Since  $B_{n-1}(\Sigma_\sigma) = \text{im } d'_n = 0$  we have

$$\tilde{H}_{n-1}(\Sigma_\sigma) = Z_{n-1}(\Sigma_\sigma).$$

Now,

$$Z_{n-1}(\Sigma_\sigma) = \ker d'_{n-1} = \ker d_{n-1} = \text{im } d_n,$$

Since  $\tilde{H}_{n-1}(K_\sigma) = 0$ .

Now  $C_n(K_\sigma) = \langle \sigma \rangle \cong \mathbb{Z}$ . Thus  $\text{im } d_n$  is generated by  $d\sigma$ . Thus  $Z_{n-1}(\Sigma_\sigma)$  is generated by  $d\sigma$  and is free by definition. Thus,

$$\tilde{H}_n(\Sigma_\sigma) = Z_n(\Sigma_\sigma) = \langle d\sigma \rangle \cong \mathbb{Z}.$$

